

## Quality Measures by Type for Optometrists

Read all the measure specifications and requirements to ensure you are properly applying each measure.

[Links to all Measure Specifications](#)

### Quality Measures: Claims

- 001 Diabetes: Glycemic Status Assessment Greater Than 9%**      **Intermediate Outcome**      **High Priority**      **Traditional MIPS**  
Percentage of patients 18-75 years of age with diabetes who had a glycemic status assessment (hemoglobin A1c [HbA1c] or glucose management indicator [GMI]) > 9.0% during the measurement period.
- 128 Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan**      **Not for Traditional MIPS-MVP only**  
Percentage of patients aged 18 years and older with a BMI documented during the current encounter or within the previous twelve months AND who had a follow-up plan documented if most recent BMI was outside of normal parameters.
- 141 Primary Open-Angle Glaucoma (POAG): Reduction of Intraocular Pressure (IOP) by 20% OR Documentation of a Plan of Care**      **Outcome**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 18 years and older with a diagnosis of primary open-angle glaucoma (POAG) whose glaucoma treatment has not failed (the most recent IOP was reduced by at least 20% from the pre-intervention level) OR if the most recent IOP was not reduced by at least 20% from the pre-intervention level, a plan of care was documented within the 12 month performance period.
- 181 Elder Maltreatment Screen and Follow-Up Plan**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 60 years and older with a documented elder maltreatment screen using an Elder Maltreatment Screening tool on the date of encounter AND a documented follow-up plan on the date of the positive screen.
- 226 Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 12 years and older who were screened for tobacco use one or more times during the measurement period AND who received tobacco cessation intervention during the measurement period or in the six months prior to the measurement period if identified as a tobacco user.
- 236 Controlling High Blood Pressure**      **Intermediate Outcome**      **High Priority**      **Traditional MIPS**  
Percentage of patients 18-85 years of age who had a diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period, and whose most recent blood pressure was adequately controlled (<140/90mmHg) during the measurement period.
- 317 Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented**      **Traditional MIPS**  
Percentage of patient visits for patients aged 18 years and older seen during the measurement period who were screened for high blood pressure AND a recommended follow-up plan is documented, as indicated, if blood pressure is elevated or hypertensive.

## Quality Measures: eCQM

- 001 Diabetes: Glycemic Status Assessment Greater Than 9%**      **Intermediate Outcome**      **High Priority**      **Traditional MIPS**  
Percentage of patients 18-75 years of age with diabetes who had a glycemic status assessment (hemoglobin A1c [HbA1c] or glucose management indicator [GMI]) > 9.0% during the measurement period.
- 012 Primary Open-Angle Glaucoma (POAG): Optic Nerve Evaluation**      **Traditional MIPS**  
Percentage of patients aged 18 years and older with a diagnosis of primary open-angle glaucoma (POAG) who have an optic nerve head evaluation during one or more visits within 12 months.
- 019 Diabetic Retinopathy: Communication with the Physician Managing Ongoing Diabetes Care**      **Traditional MIPS**  
Percentage of patients aged 18 years and older with a diagnosis of diabetic retinopathy who had a dilated macular or fundus exam performed with documented communication to the physician who manages the ongoing care of the patient with diabetes mellitus regarding the findings of the macular or fundus exam at least once during the performance period.
- 117 Diabetes: Eye Exam**      **Traditional MIPS**  
Percentage of patients 18-75 years of age with diabetes and an active diagnosis of retinopathy in any part of the measurement period who had a retinal or dilated eye exam by an eye care professional during the measurement period or diabetics with no diagnosis of retinopathy in any part of the measurement period who had a retinal or dilated eye exam by an eye care professional during the measurement period or in the 12 months prior to the measurement period.
- 128 Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan**      **Not for Traditional MIPS-MVP only**  
Percentage of patients aged 18 years and older with a BMI documented during the current encounter or within the previous twelve months AND who had a follow-up plan documented if most recent BMI was outside of normal parameters.
- 130 Documentation of Current Medications in the Medical Record**      **High Priority**      **Traditional MIPS**  
Percentage of visits for which the eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter.
- 226 Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 12 years and older who were screened for tobacco use one or more times during the measurement period AND who received tobacco cessation intervention during the measurement period or in the six months prior to the measurement period if identified as a tobacco user.
- 236 Controlling High Blood Pressure**      **Intermediate Outcome**      **High Priority**      **Traditional MIPS**  
Percentage of patients 18-85 years of age who had a diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period, and whose most recent blood pressure was adequately controlled (<140/90mmHg) during the measurement period.
- 238 Use of High-Risk Medications in Older Adults**      **High Priority**      **Traditional MIPS**  
Percentage of patients 65 years of age and older who were ordered at least two high-risk medications from the same drug class.
- 281 Dementia: Cognitive Assessment**      **Traditional MIPS**  
Percentage of patients, regardless of age, with a diagnosis of dementia for whom an assessment of cognition is performed and the results reviewed at least once within a 12-month period.
- 317 Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented**      **Traditional MIPS**  
Percentage of patient visits for patients aged 18 years and older seen during the measurement period who were screened for high blood pressure AND a recommended follow-up plan is documented, as indicated, if blood pressure is elevated or hypertensive.
- 318 Falls: Screening for Future Fall Risk**      **High Priority**      **Traditional MIPS**  
Percentage of patients 65 years of age and older who were screened for future fall risk during the measurement period.
- 374 Closing the Referral Loop: Receipt of Specialist Report**      **High Priority**      **Traditional MIPS**  
Percentage of patients with referrals, regardless of age, for which the referring clinician receives a report from the clinician to whom the patient was referred.

## Quality Measures: CQM

- 001 Diabetes: Glycemic Status Assessment Greater Than 9%**      **Intermediate Outcome High Priority**      **Traditional MIPS**  
Percentage of patients 18-75 years of age with diabetes who had a glycemic status assessment (hemoglobin A1c [HbA1c] or glucose management indicator [GMI]) > 9.0% during the measurement period.
- 117 Diabetes: Eye Exam**      **Traditional MIPS**  
Percentage of patients 18-75 years of age with diabetes and an active diagnosis of retinopathy in any part of the measurement period who had a retinal or dilated eye exam by an eye care professional during the measurement period or diabetics with no diagnosis of retinopathy in any part of the measurement period who had a retinal or dilated eye exam by an eye care professional during the measurement period or in the 12 months prior to the measurement period.
- 128 Preventive Care and Screening: Body Mass Index (BMI) Screening and Follow-Up Plan**      **Not for Traditional MIPS-MVP only**  
Percentage of patients aged 18 years and older with a BMI documented during the current encounter or within the previous twelve months AND who had a follow-up plan documented if most recent BMI was outside of normal parameters.
- 130 Documentation of Current Medications in the Medical Record**      **High Priority**      **Traditional MIPS**  
Percentage of visits for which the eligible clinician attests to documenting a list of current medications using all immediate resources available on the date of the encounter.
- 141 Primary Open-Angle Glaucoma (POAG): Reduction of Intraocular Pressure (IOP) by 20% OR Documentation of a Plan of Care**      **Outcome High Priority**      **Traditional MIPS**  
Percentage of patients aged 18 years and older with a diagnosis of primary open-angle glaucoma (POAG) whose glaucoma treatment has not failed (the most recent IOP was reduced by at least 20% from the pre-intervention level) OR if the most recent IOP was not reduced by at least 20% from the pre-intervention level, a plan of care was documented within the 12 month performance period.
- 155 Falls: Plan of Care**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 65 years and older with a history of falls that had a plan of care for falls documented within 12 months.
- 181 Elder Maltreatment Screen and Follow-Up Plan**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 60 years and older with a documented elder maltreatment screen using an Elder Maltreatment Screening tool on the date of encounter AND a documented follow-up plan on the date of the positive screen.
- 226 Preventive Care and Screening: Tobacco Use: Screening and Cessation Intervention**      **High Priority**      **Traditional MIPS**  
Percentage of patients aged 12 years and older who were screened for tobacco use one or more times during the measurement period AND who received tobacco cessation intervention during the measurement period or in the six months prior to the measurement period if identified as a tobacco user.
- 236 Controlling High Blood Pressure**      **Intermediate Outcome High Priority**      **Traditional MIPS**  
Percentage of patients 18-85 years of age who had a diagnosis of essential hypertension starting before and continuing into, or starting during the first six months of the measurement period, and whose most recent blood pressure was adequately controlled (<140/90mmHg) during the measurement period.
- 238 Use of High-Risk Medications in Older Adults**      **High Priority**      **Traditional MIPS**  
Percentage of patients 65 years of age and older who were ordered at least two high-risk medications from the same drug class.
- 282 Dementia: Functional Status Assessment**      **Traditional MIPS**  
Percentage of patients with dementia for whom an assessment of functional status was performed at least once in the last 12 months.
- 286 Dementia: Safety Concern Screening and Follow-Up for Patients with Dementia**      **High Priority**      **Traditional MIPS**  
Percentage of patients with dementia or their caregiver(s) for whom there was a documented safety concerns screening in two domains of risk: 1) dangerousness to self or others and 2) environmental risks; and if safety concerns screening was positive in the last 12 months, there was documentation of mitigation recommendations, including but not limited to referral to other resources.

- 288 Dementia: Education and Support of Caregivers for Patients with Dementia High Priority Traditional MIPS**  
Percentage of patients with dementia whose caregiver(s) were provided with education on dementia disease management and health behavior changes AND were referred to additional resources for support in the last 12 months.
- 317 Preventive Care and Screening: Screening for High Blood Pressure and Follow-Up Documented Traditional MIPS**  
Percentage of patient visits for patients aged 18 years and older seen during the measurement period who were screened for high blood pressure AND a recommended follow-up plan is documented, as indicated, if blood pressure is elevated or hypertensive.
- 374 Closing the Referral Loop: Receipt of Specialist Report High Priority Traditional MIPS**  
Percentage of patients with referrals, regardless of age, for which the referring clinician receives a report from the clinician to whom the patient was referred.
- 431 Preventive Care and Screening: Unhealthy Alcohol Use: Screening & Brief Counseling Traditional MIPS**  
Percentage of patients aged 18 years and older who were screened for unhealthy alcohol use using a systematic screening method at least once within the last 12 months AND who received brief counseling if identified as an unhealthy alcohol user.
- 487 Screening for Social Drivers of Health High Priority Traditional MIPS**  
Percent of patients 18 years and older screened for food insecurity, housing instability, transportation needs, utility difficulties, and interpersonal safety.
- 489 Connection to Community Service Provider High Priority Traditional MIPS**  
Percent of patients 18 years or older who screen positive for one or more of the following health related social needs (HRSNs): food insecurity, housing instability, transportation needs, utility help needs, or interpersonal safety; and had contact with a Community Service Provider (CSP) for at least 1 of their HRSNs within 60 days after screening.
- 499 Appropriate screening and plan of care for elevated intraocular pressure following intravitreal or periocular steroid therapy Traditional MIPS**  
Percentage of patients who had an intravitreal or periocular corticosteroid injection (e.g., triamcinolone, preservative-free triamcinolone, dexamethasone, dexamethasone intravitreal implant, or fluocinolone intravitreal implant) who, within seven (7) weeks following the date of injection, are screened for elevated intraocular pressure (IOP) with tonometry with documented IOP  $\leq$ 25 mm Hg for injected eye OR if the IOP was  $>$ 25 mm Hg, a plan of care was documented.
- 500 Acute posterior vitreous detachment appropriate examination and follow-up Traditional MIPS**  
Percentage of patients with a diagnosis of acute posterior vitreous detachment (PVD) in either eye who were appropriately evaluated during the initial exam and were re-evaluated no later than 8 weeks.
- 501 Acute posterior vitreous detachment and acute vitreous hemorrhage appropriate examination and follow-up Traditional MIPS**  
Percentage of patients with a diagnosis of acute posterior vitreous detachment (PVD) and acute vitreous hemorrhage in either eye who were appropriately evaluated during the initial exam and were re-evaluated no later than 2 weeks.
- 508 Adult COVID-19 Vaccination Status Traditional MIPS**  
Percentage of patients aged 18 years and older seen for a visit during the performance period that are up to date on their COVID-19 vaccinations as defined by Centers for Disease Control and Prevention (CDC) recommendations on current vaccination.

#### Cannot be used with Post-Operative Modifiers

- 191 Cataracts: 20/40 or Better Visual Acuity within 90 Days Following Cataract Surgery** (eCQM, CQM Outcome High Priority Traditional MIPS)
- 303 Cataracts: Improvement in Patient's Visual Function within 90 Days Following Cataract Surgery** (CQM-Patient reported Outcome-High Priority Traditional MIPS)
- 304 Cataracts: Patient Satisfaction within 90 Days Following Cataract Surgery** (CQM, High Priority Traditional MIPS)
- 384 Adult Primary Rhegmatogenous Retinal Detachment Surgery: No Return to the Operating Room Within 90 Days of Surgery** (CQM Outcome-High Priority Traditional MIPS)
- 385 Adult Primary Rhegmatogenous Retinal Detachment Surgery: Visual Acuity Improvement Within 90 Days of Surgery** (CQM Outcome-High Priority Traditional MIPS)
- 389 Cataract Surgery: Difference Between Planned and Final Refraction** (CQM Outcome-High Priority Traditional MIPS)